


# FIG. 1


## INFLUENCE OF FEED ON AVERAGE WEIGHT (BWT), FEED CONVERSION (FC), ADJUSTED FEED CONVERSION (AFC), AND MORTALITY

COMPOSITION		DAY 18			DAY 50			7-49 DAYS
NO	CITRUS PEEL LBS/TON	BWT (KG)	FC (KG GAIN/ KG FEED)	AFC (KG GAIN KG FEED)	BWT (KG)	FC (KG GAIN/ KG FEED)	AFC (KG GAIN/ KG FEED)	MORTALITY (%)
1	CONTROL	0.583±0.007	1.252±0.007	1.243±0.007	2.860±0.023	1.987±0.018	1.923±0.014	6.000±1.309
2	4	0.582±0.006	1.242±0.003	1.231±0.002	2.793±0.025	1.964±0.014	1.915±0.009	4.250±0.796
3	8	0.581±0.005	1.233±0.007	1.220±0.007	2.824±0.022	1.953±0.016	1.910±0.010	3.250±0.750
4	16	0.582±0.007	1.244±0.009	1.237±0.009	2.835±0.024	1.959±0.014	1.912±0.011	3.750±1.098
5	24	0.585±0.006	1.249±0.006	1.241±0.007	2.853±0.022	1.970±0.018	1.919±0.006	4.250±1.278
6	32	0.577±0.008	1.259±0.009	1.245±0.008	2.890±0.023	2.000±0.025	1.910±0.010	7.250±1.359
P<F		0.9631	0.1862	0.1262	0.0916	0.4288	0.9335	0.1288

# FIG.2




# FIG.3




# FIG.4

TOTAL MORTALITY,  
SDS MORTALITY AND ACT/ACT-S MORTALITY




# FIG.5


CUMULATIVE MORTALITY (%)



# FIG. 6




# FIG. 7



# FIG.8

AVERAGE BODY WEIGHTS, GAIN  
AND ADJUSTED FEED CONVERSION  
(DAYS 0-15)



# FIG.9

AVERAGE BODY WEIGHT DAY 27 AND  
ADJUSTED FEED CONVERSION

